

Quadratic Optimization Problem on Permutation Set with Simulation of Applied Tasks

Liudmyla Koliechkina¹[0000-0002-4079-1201], Alla Nahirna²[0000-0002-2863-8706],

Olena Dvirna³[0000-0002-0750-6958]

¹University of Lodz, Uniwersytecka Str. 3, 90-137 Lodz, Poland
ludaplt1971@gmail.com

²Kyiv International University, Lvivska str., 49, Kyiv, 03179, Ukraine
naghirnaalla@ukr.net

³Poltava University of Economics and Trade, Koval str., 3, Poltava, 36000, Ukraine
lenadvirna@gmail.com

Abstract. The article discusses the formulation of an optimization problem with a quadratic target function and additional constraints on the permutation set, which can be a model of many applied problems. An algorithm for solving an optimization problem with a quadratic target function and additional constraints on permutations is proposed.

During the implementation of the method the first reference plan is found and additional restrictions for it are checked at the first stage. Thus, in the beginning of the algorithm, the number of considered solutions decreases. This makes it possible at the first stage to reduce the number of possible solutions and narrow the area of the problem study. An example of solving a theoretical problem using this method, demonstrating its effectiveness, is proposed.

Such task can be used to modeling various technological processes. The reason for this is the optimization of mathematical models and algorithms for the proposed models.

Keywords: optimization problems, combinatorial set of permutations, model of optimization problems, quadratic target function, optimal solutions.

1 Introduction

Computer simulation of technological processes consists in optimization of the study object by a mathematical model, and further model study with the help of implemented computational algorithms on personal computers. The computer simulation process, as a single process of building and researching a model, is used for research, analysis, design and optimization of technological objects and systems [1]. Today, computer modeling approaches combines both theory and practice. Working with a

model that represents a research object gives you the opportunity to explore its properties and behavior in all situations.

Computer modeling is the process of building a real study object's model and setting up computational experiments on this model in order to understand and evaluate various strategies based on the use of algorithms that ensure the functioning of this study object. Thus, the process of computer simulation includes the model design, and its application to solve the problem of optimization and design of technological processes in production. All these problems are extremely complex and include numerous elements, variables, parameters, constraints, etc. Such problems can be modeled by combinatorial optimization tasks [1-8].

The combinatorial optimization problems study comprises a fairly wide range of mathematical models associated with the need to solve various important practical problems of optimal planning, management and design [1, 2, 5, 7, 9, 10-12, 20-29]. In this regard, many papers have recently appeared which investigate the combinatorial optimization problems and propose approaches to their solution [8, 11-34, 36-38].

The paper presents a model of optimization problems on a combinatorial set of permutations, which is a model of many applied problems. It is proposed the algorithm for solving an optimization problem with a quadratic target function and additional restrictions on permutations.

During the method's implementation the first reference plan is located and additional restrictions are checked, which reduces the number of considered solutions at the beginning of the algorithm

It is offered an example of solving a theoretical problem by the given method and demonstrating its efficiency.

2 Literature review

Computer models have become a common tool for mathematical modeling and are used in various areas of electronics, engineering, industry, and so on. Computer models are used to obtain new knowledge about the object or for an approximate assessment of the system's behavior that are too complex for analytical research.

Computer modeling is one of the effective methods for studying complex systems. The construction of a computer model is based on abstracting from the specific nature of the phenomena or the original object under study and consists of two stages – first, the creation of a qualitative, and then a quantitative model. The more significant properties will be identified and transferred to the computer model – the more approximate it will be to the real model, the greater the capabilities that the system using this model will have. Today the works of many scientists are devoted to this research area [1-5, 8-10, 21].

An important computer simulation area is analytical and simulation modeling. In analytical modeling, mathematical (abstract) models of a real object are studied in the algebraic equations form, as well as providing for the implementation of an unambiguous computational procedure leading to their exact solution. In simulation, mathematical models are studied in the algorithms form, which reproduce the

functioning of the system under study by sequentially performing numerous elementary operations. Very relevant today for simulation and analytical modeling are discrete models, in particular, combinatorial optimization, their study is the subject of a great number of papers [1–17], [22–31].

A significant contribution to the development of modern discrete optimization theory, the development and implementation of its methods in the study and solving of important applied problems have been made over the course of more than forty years under the scientific guidance of Academician I.V. Sergienko scientists of the Institute of Cybernetics named by V.M. Glushkov National Academy of Sciences of Ukraine. They obtained important results, which became the basis for the theory development and the creation of new lines in discrete optimization.

At present, intensive studies of the stability problems of vector discrete optimization problems are carried out at the Institute of Cybernetics named by V.M. Glushkov of the National Academy of Sciences of Ukraine (I.V. Sergienko, T.T. Lebedev, N.V. Semenova, T.I. Sergienko) [16, 17], Belarusian State University (V.O. Emelyichev, D.P. Podkopyayev, Yu.V. Nikulin, etc.) [6, 7], the Joint Institute of Computer Sciences of the National Academy of Sciences of Belarus (Yu.N. Sotskov), the Computational Center of the Russian Academy of Sciences (E.M. Gordeyev, V.K. Leontiev), Omsk Branch of the Institute of System Studies of the Polish Academy of Sciences (M. Libura), a number of universities of the Federal Republic of Germany (E. Girlich), the Netherlands (ES van der Poort, G. Sierkma, APM Wagelmans, JAA van der Veen), in the uniquely Colorado US (H.J. Greenberg).

Belarusian scientists under the scientific guidance of Professor V.O. Yemelyicheva successfully develops a constructive approach to the problem of the vector discrete optimization problems correctness associated with the reception of quantitative characteristics of the stability of specified problem statements, a mathematical apparatus for studying the stability of multicriteria discrete problems with different types, vector criteria functions, principles of optimality, as well as generalized and parameterized principles of optimality are developed.

It is important to study the relation between different classes of point configurations triangulation (regular, weakly regular, deployed, symmetric, political, etc.), in particular, the study of the structure of a partially ordered set constructed on the family of triangulation classes considered in relation to the inclusion relation. Recently, algorithms for solving convex polyhedral admissibility problems are actively explored [11, 18, 19]. One of the interesting computational approaches is the reduction of the admissibility problem for a polyhedron to the projection problem on a normal cone generated by a dual system of inequalities [18, 29, 31], which is sufficiently close to the projection problem for a binary polyhedron. Today, in the research area of various classes of combinatorial models, the new methods development for their solution, great attention is paid to methods based on the use of structural properties of combinatorial sets. The properties of combinatorial sets study is closely related to the theory of polyhedral and graphs. The use of information about the structure of the convex shell of admissible multivariate solutions, which is the basis for many methods, is one of the most successful approaches to solving combinatorial optimization problems for today. But when solving such tasks there are

problems related to the complexity of mathematical models, large volume of information, etc.

Applied problems simulated by extreme discrete tasks often have a high dimensionality, so they are quite complex from a computational point of view. The main task is to determine the value of an argument belonging to a certain combinatorial configuration for which the target function acquires the global optimum. So it is necessary to develop the most effective algorithm, which is based on the specific properties of the combinatorial configurations.

For the extreme problems of combinatorial optimization, polynomial algorithms for finding an optimal solution based on the properties of the input data structure have been developed, but there are few such work compared to methods based on partial overview of the options. One of the approaches to solving such problem is to carry out research and analysis of the combinatorial configurations properties, in which the target function, which reflects the combinatorial nature of the tasks, is determined. Analysis and study of combinatorial configurations as a target function argument, setting the change in the values of the target function, depending on the elements ordering of the selected combinatorial configuration and the structure of the input data specificity, does not pay sufficient attention in the literature. But it should be noted that the study of the certain tasks properties in order to identify their characteristic properties and their use for solving the problem, gives the possibility of constructing new approaches and the development of known methods.

Hence, one of the important problems in the discrete optimization area is the detection of the properties of combinatorial configurations in extreme problems, the use of which would allow to establish the regularity of the change in the values of the target functions, depending on the argument ordering and on the specificity and structure of the combinatorial configurations sets.

Today, significant results have been obtained in the area of research of combinatorial models' various classes and the development of new methods for their solution. The following foreign scientists made a fundamental contribution to the development of discrete, in particular, combinatorial optimization: M. Gary, S. Berge, D. Johnson, H. Papadimitriou, P. Pardalos, K. Staiglich, R. Stanley, F. Harari, V.A. Emelichev, V.M. Sachkov [1-8, 10, 11, 12].

In turn, the many Ukrainian scientists' works are devoted to the various classes of combinatorial optimization problems' study: L.F. Gulyanitsky, P.I. Stetsyuka, I.V. Sergienko, N.S. Shor, Yu.G. Stoyan, S.V. Yakovlev, A.O. Yemetsa, V.O. Perepelitsy and many others.

In particular, in [18–21], the authors describe the convex extensions theory and its applications in combinatorial optimization problems. Combinatorial models' applications in practical problems of geometric design are presented in the works of L.F. Gulyanitsky, I.V. Sergienko, Yu.G. Stoyan, S.V. Yakovlev, N.S. Shor [22-27, 38].

Quadratic optimization on the permutation set is reflected in [28–31].

The permutation set's representation as the intersection of a permutation polyhedron and a hypersphere is interesting, as well as optimization methods on permutation configurations using the intersection described in [32].

The development of an integrated approach to the analysis of the properties of combinatorial optimization supplies covers a wide range of studies of combinatorial functions, combinatorial polyhedral, combinatorial configurations as an argument of the target function. The results give the opportunity to improve existing methods for solving such problems and develop new methods for optimal solutions searching. The problems based on the properties of combinatorial configurations are actual problems of combinatorial optimization. Of particular importance in this aspect is the consideration of extreme problems in combinatorial configurations using graph theory.

The research of tasks in graphs deals with such scientists as F. Harari, O. Ore, I.V. Sergienko, V. O. Yemelichev, A. O. Zikov, V. O. Perepelytsya, R. I. Tyshkevich and others. Despite quite large achievements in the area of discrete optimization, in the process of modeling, there are extreme problems classes for which a number of issues have not yet been investigated. Principal difficulties that arise during modeling are also related to various types of uncertainty. These include: the availability of many criteria for evaluating the quality of solutions, interval setting of task parameters, etc. In these conditions, classical methods are not sufficiently suitable for solving problems. As you know, most combinatorial optimization tasks can be reduced to integer programming tasks, but this is not always justified, since it eliminates the possibility of taking into account the combinatorial properties of task solutions.

This work is a continuation of research in the extreme discrete problem's area, in particular, combinatorial optimization, as well as optimization problems under the conditions of multicriteria, which arise in the study of many theoretical and applied problems.

3 Formal problem statement

We consider the permutation as an ordered sample of elements $a = (a_{i_1}, a_{i_2}, \dots, a_{i_k})$, where $a_{i_j} \in A$, $\forall i_j \in N_n$, $\forall j \in N_n$ $i_s \neq i_t$ if $s \neq t \forall s \in N_n$, $\forall t \in N_n$ from some multi-set $A = \{a_1, a_2, \dots, a_q\}$, which is characterized by the base $S(A) = \{e_1, e_2, \dots, e_k\}$, where $e_j \in R^1$, $\forall j \in N_k$ and the multiplicity of the elements $k(e_j) = r_j$, $j \in N_k$, $r_1 + r_2 + \dots + r_k = q$ $j \in N_k$, according to [14, 17].

A set of permutations with repetitions of n real numbers, among which k different, is called the general permutation set and is denoted as $P_{nk}(A)$. This is the set of ordered n -samples from the multiset A under the condition $n = q > k$.

If we have $n = k = q$ set of permutations without repetition, we denote it as P_n . Obviously, $P_n(A) = P_{nn}(A)$. In cases where the form of the set of permutations is not indicated, it will be possible to write down these sets as $P(A)$.

It is known [14] that the convex hull of the set of permutations is a permutation polyhedron $\Pi = \text{conv } P(A)$, which vertex set $P(A)$ is equal to the set of permutations: $\text{vert } \Pi(A) = P(A)$.

Without loss of generality, we order the elements of the multiset A in non-descending order: $a_1 \leq a_2 \leq \dots \leq a_q$ and the elements of its foundation—in descending order: $e_1 > e_2 > \dots > e_k$.

Then the convex hull of a common set of $P(A)$ permutations is a generic polyhedron $\Pi(A) = \text{conv } P(A)$, that is described by a well-known system of linear inequalities:

$$\begin{cases} \sum_{j=1}^n x_j \leq \sum_{j=1}^n a_j, \\ \sum_{j=1}^i x_{\alpha_j} \leq \sum_{j=1}^i a_j, \end{cases} \quad (1)$$

$\alpha_j \in N_n, \alpha_j \neq \alpha_t, \forall j \neq t, \forall j, t \in N_i, \forall i \in N_n$ and $P(A) = \text{vert } \Pi(A)$.

Consider the optimization problem:

$$Z(\Phi, P(A)) : \max \{\Phi(a) \mid a \in P(A)\} \quad (2)$$

$$D = \{x \in R^n \mid Gx \leq (\geq) b\} \quad (3)$$

where $G \in R^{m \times n}$, $b \in R^m$

$$\Phi(a) = \sum_{j=1}^n c_j x_j^2 + \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j \quad (4)$$

on a set of permutations $P_n(A)$.

Additional linear constraints form a multifaceted set $D \subset R^n$.

Then, to every point $a \in P_n(A)$ will match point $x \in X$, one that satisfies equality $F(x) = \Phi(a)$:

$$Z(F, X) : \max \{F(x) \mid x \in X\} \quad (5)$$

$$\text{where, } F(x) = \sum_{j=1}^n c_j x_j^2 + \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j \quad (6)$$

and additional constraints:

$$D = \{x \in R^n \mid Gx \leq (\geq) b\} \quad (7)$$

where $G \in R^{m \times n}$, $b \in R^m$

X - nonempty set in R^n , which is defined as follows:

$$X = \text{vert } \Pi(A), \quad \Pi = \text{conv } P(A) \quad (8)$$

It is natural to assume that the maximum of a quadratic function will be one of the vertices of the permutation polyhedron, and the vertices of the graph $G(P_n)$ will match to all points of the set of permutations $P_n(A)$.

The adjacency of the vertices of a permutation polyhedron is determined by a one-time transposition of two vertex elements. The number of transpositions in the graph is determined by the formula [35]:

$$C_n^2 = \frac{n(n-1)}{2} \quad (9)$$

4 The algorithm for solving an optimization problem with a quadratic target function and additional constraints on the set is rearranged

The algorithm for finding the problem optimal solution consists of four steps, which in a few steps make it possible to obtain an optimal solution.

STEP 1. Finding the first support solution.

Consider the first transposition of the target function:

$$x_1 \leftrightarrow x_2, (i = 2, \dots, n):$$

$$f_{12} = (x_1 - x_2)(a_1x_1 + a_1x_2 + \dots + a_nx_n)$$

We form the necessary conditions for finding the first solution:

$$\begin{cases} x_1 > x_2, \\ x_i \rightarrow \max, (\max a_i). \end{cases} \quad (10)$$

Variants of transpositions:

$$x_1 \leftrightarrow x_2, (i = 2, \dots, n):$$

$$f_{12} = (x_1 - x_2)(a_1x_1 + a_1x_2 + \dots + a_nx_n)$$

$$x_1 \leftrightarrow x_3 : f_{13} = (x_1 - x_3)(a_{13}x_1 + a_{23}x_2 + \dots + a_{n3}x_n)$$

... ..

$$x_1 \leftrightarrow x_n : f_{1n} = (x_1 - x_n)(a_{1n}x_1 + a_{2n}x_2 + \dots + a_{nn}x_n) \quad (11)$$

$$x_2 \leftrightarrow x_i, (i = 3, \dots, n);$$

• • • • •

• • • • •

The first solution will be: (x_1, x_2, \dots, x_n) . It should be noted that there may be several first solutions.

$$\Delta g_i = \Delta g_i^2 - \Delta g_i^1 = \left(x_i^{g_i^2} * c_j + x_j^{g_i^2} * c_i \right) - \left(x_j^{g_i^1} * c_j + x_i^{g_i^1} * c_i \right) \quad (14)$$

All increments satisfy conditions (12), then the found point is the optimal solution. Otherwise, we return to the beginning of the step 4.

4.2. If condition (13) is not fulfilled, we return to the beginning of step 4.

It should be noted that conditions (12) are sufficient for finding the optimal solution, and the fulfillment of inequality (13) is necessary for finding the optimal solution.

5 Example

Find the maximum value of the target function:

$$\max F(x) = \frac{1}{2}(2x_1 - x_2 - x_3 + x_4)^2 + (x_2 + x_3)^2 - \frac{1}{2}(x_3 - x_2 + 2x_4)^2 + 2x_4^2$$

on the permutations set A_4 , where $A = (1,2,3,4)$, with the following restrictions:

$$\begin{cases} g_1 = 5x_1 - 2x_2 + 3x_3 + 4x_4 \geq 15, \\ g_2 = -3x_1 + 6x_2 + 8x_3 - x_4 \leq 31. \end{cases}$$

Solving.

For formula (1), the number of possible transpositions is: $C_4^2 = 6$.

Consider the first transposition $x_1 \leftrightarrow x_2$, presenting the target function as a product:

$$f_{12} = (x_1 - x_2)(x_1 + x_2 - 6x_3 + x_4),$$

accordingly, when searching for the first solution, it is necessary to consider the following conditions:

$$\begin{cases} x_1 > x_2, \\ x_3 \rightarrow \min. \end{cases}$$

Then the first solution may be the following points of the permutations set: (4,3,1,2), (4,2,1,3), (3,2,1,4).

Consider the first point (4,3,1,2):

$$\begin{cases} g_1(4,3,1,2) = 25 \geq 15, \\ g_2(4,3,1,2) = 12 \leq 31. \end{cases}$$

Additional restrictions are enforced. Then the initial conditions for improving the first support solution will be:

$$\max f_1(4,3,1,2) = 40,$$

$$\begin{cases} \Delta g_1^1(4,3,1,2) \geq -10, \\ \Delta g_2^1(4,3,1,2) \leq 19. \end{cases}$$

When considering point 2, it is necessary to fulfill the condition Δf_i - increases:

$$\Delta f_2(4,2,1,3) = (x_2 - x_4)(-4x_1 + 0.5x_2 + 0.5x_4 + 7x_3) = 6,5.$$

Consequently, the target function increases by 6.5 units, so there is a need to check the increment of additional restrictions:

$$\begin{cases} \Delta g_1^1(4,2,1,3) = 6 \geq -10, \\ \Delta g_2^1(4,2,1,3) = -7 \leq 19. \end{cases}$$

Constraints are met, so the support solution can be improved:

$$\max f_2(4,2,1,3) = f_1(4,3,1,2) + \Delta f_2(4,2,1,3) = 46,5.$$

$$\begin{cases} \Delta g_1^1(4,2,1,3) \geq -16, \\ \Delta g_2^1(4,2,1,3) \leq 26. \end{cases}$$

Consider the point (3,2,1,4), then you need to transpose $x_1 \leftrightarrow x_4$ regarding point 2 (4,2,1,3) :

$$\Delta f_{14}(3,2,1,4) = \frac{1}{2}(x_1 - x_4)(3x_1 - 6x_2 + 2x_3 + 3x_4) = \frac{1}{2}(3 - 4)(9 - 12 + 2 + 12) = -5,5.$$

Since the target function decreases by 5.5 units, there is no point in considering this point of the set of permutations.

Consider ascending, point (4,2,3,1), respectively, transposition, $x_3 \leftrightarrow x_4$ regarding point 2 (4,2,1,3) :

$$\Delta f_{34}(3,2,1,4) = \frac{1}{2}(x_3 - x_4)(6x_2 - 8x_1 + x_3 + x_4) = \frac{1}{2}(3 - 1)(12 - 32 + 3 + 1) = -16.$$

The target function decreases by 16 units, respectively, this point is not considered.

Therefore, point 2 (4,2,1,3) is optimal and cannot be improved.

$$\max f_2(4,2,1,3) = 46,5.$$

6 Conclusion

The article represents an optimization problem model with a quadratic target function and additional constraints on the combinatorial set of permutations as a model of many applied problems. An algorithm for solving the optimization problem is proposed and a numerical example is demonstrated. It should be noted that the problem are very complex from a computational point of view. Therefore, their solutions re-

quire a lot of time and resources. This method significantly simplifies the procedure for finding the optimal solution of an optimization problem with a quadratic target function and additional constraints on the combinatorial set of permutations, since the inequality of restrictions increments allows you to immediately determine whether the point of the permutation set is a support solution or not. There is not a necessity to do complex calculations of all constraints and the target function; it suffices to find the increment of constraint and functions in the case of an improvement the solution.

The further development of this study is going to be aimed at realizing and adapting the formulated method on other combinatorial constructions, as well as developing new methods for solving combinatorial optimization problems, taking into account the input data.

References

1. Korte, B., Vygen, J.: Combinatorial Optimization: Theory and Algorithms. Springer, New York (2018)
2. Pardalos, P.M., Du, D.-Z., Graham, R.L. (Eds.): Handbook of combinatorial optimization. Springer, New York (2013)
3. Bernhard, K., Jens, V.: Combinatorial Optimization, Theory and Algorithms. Springer-Verlag, Berlin (2012)
4. Karin, R., Saoub, A.: Tour through Graph Theory. Chapman and Hall/CRC, New York (2017)
5. Berge, C.: Principes de combinatoire. Dunod, Paris (1968)
6. Yemelichev, V.A., Kovalev, M.M., Kravtsov, M.K.: Polytopes, graphs and optimisation. Cambridge University Press, Cambridge (1984)
7. Sachkov, V.N.: Combinatorial methods of discrete mathematics. Nauka, Moscow (1975)
8. Hulianytskyi, L., Riasna, I.: Formalization and classification of combinatorial optimization problems. Springer Optimization and its Applications. 130: pp. 239-250 (2017)
9. Yakovlev, S.V., Grebennik, I.V.: Localization of solutions of some problems of nonlinear integer optimization. Cybernetics and Systems Analysis. 29(5): pp. 727-734 (1993)
10. Schrijver, A.: Combinatorial Optimization: Polyhedra and Efficiency. Springer Science & Business Media, Berlin (2012)
11. Balakrishnan, R., Sridharan, S.: Foundations of Discrete Mathematics with Algorithms and Programming. Chapman and Hall/CRC, New York (2018)
12. Bona, M.: Combinatorics of Permutations, Second Edition. Chapman and Hall/CRC, New York (2012)
13. Shor, N.Z., Stetsyuk, P.I.: Modified r-algorithm to find the global minimum of polynomial functions Cybernetics and Systems Analysis. 33(4): pp.482-497 (1997)
14. Emets, O.A., Koliechkina, L. N.: Solution of optimization problems with fractional-linear objective functions and additional linear constraints on permutations. Cybernetics and Systems Analysis. 40(3): pp. 329-339 (2004)
15. Donec, G. A., Kolechkina, L. M.: Method of ordering the values of a linear function on a set of permutations. Cybernetics and Systems Analysis. 45(2): 204-213 (2009)
16. Sergienko, I.V., Volkovich, O.V., Roshchin, V.A.: Models and methods of solution of quadratic integer programming problems. Cybernetics. 23(3): 289-305 (1987)

17. Semenova, N.V., Kolechkina, L.N., Nagornaya, A.N.: On Approach to Solving Vector Problems with Fractionally Linear Functions of the Criteria on the Combinatorial Set of Arrangements. *Journal of Automation and Information Sciences*. 42(1): pp. 67-80 (2010)
18. Yakovlev, S.V.: The theory of convex continuations of functions on vertices of convex polygons. *Computational Mathematics and Mathematical Physics*. 34: pp. 959-965 (1994)
19. Yakovlev, S.: Convex extensions in combinatorial optimization and their applications. *Springer Optimization and its Applications*. 130: pp. 567-584 (2017)
20. Yakovlev, S.V.: Bounds on the minimum of convex functions on Euclidean combinatorial sets. *Cybernetics*. 25(3): pp. 385-391 (1989)
21. Pichugina, O.S., Yakovlev, S.V.: Continuous representations and functional extensions in combinatorial optimization. *Cybernetics and Systems Analysis*. 52(6): pp. 921-930 (2016)
22. Yakovlev, S.V.: The method of artificial space dilation in problems of optimal packing of geometric objects. *Cybernetics and Systems Analysis*. 53(5): pp. 725-732 (2017)
23. Stoyan, Y.G., Yakovlev, S.V.: Configuration space of geometric objects. *Cybernetics and Systems Analysis*. 54(5): 716-726 (2018)
24. Yakovlev, S.V.: On some classes of spatial configurations of geometric objects and their formalization. *Journal of Autom and Information Sciences* . 50(9): 38-50 (2018)
25. Stoyan, Yu.G., Sokolovskii, V.Z., Yakovlev, S.V.: Method of balancing rotating discretely distributed masses. *Energomashinostroenie*. 2: pp. 4-5 (1982)
26. Stoyan, Y., Romanova, T., Pankratov, A., Kovalenko, A., Stetsyuk, P.: Balance layout problems: mathematical modeling and nonlinear optimization. *Springer Optimization and its Applications*. vol. 114, pp. 369-400 (2016)
27. Gulianitsky, L.F. Sergienko, I.V.: Meta-evolutionary method of deformed polyhedron in combinatorial optimization. *Cybernetics and system analysis*. 44(6): pp. 70-79 (2007)
28. Stoyan, Y.G., Yakovlev, S.V., Parshin, O.V.: Quadratic optimization on combinatorial sets in R^n . *Cybernetics and Systems Analysis*. 27(4): pp. 561–567 (1991)
29. Yakovlev, S.V., Valuiskaya, O.A.: Optimization of linear functions at the vertices of a permutation polyhedron with additional linear constraints. *Ukrainian Mathematical Journal*. 53(9): pp. 1535-1545 (2001)
30. Pichugina, O.S., Yakovlev, S.V.: Functional and analytic representations of the general permutations. *Eastern-European Journal of Enterprise Technologies*. 1(4): pp. 27-38 (2016)
31. Pichugina, O., Yakovlev, S.: Convex extensions and continuous functional representations in optimization, with their applications. *Journal of Coupled Systems and Multiscale Dynamics*. 4(2): pp. 129–152 (2016)
32. Yakovlev, S.V., Pichugina, O.S.: Properties of combinatorial optimization problems over polyhedral-spherical sets. *Cybernetics and Systems Analysis*. 54(1): pp. 99-109 (2018)
33. Kolichekina, L. N., Dvirna, O. A., Nagornaya, A. N.: Modified Coordinate Method to Solve Multicriteria Optimization Problems on Combinatorial Configurations. *Cybernetics and Systems Analysis*. 59(4): pp. 620-626 (2014)
34. Kolichekina, L. N., Dvirna, O. A.: Solving Extremum Problems with Linear Fractional Objective Functions on the Combinatorial Configuration of Permutations Under Multicrteriality. *Cybernetics and Systems Analysis*. 53(4): pp. 590-599 (2017)
35. Donec, G. A., Kolechkina, L. M.: Construction of hamiltonian paths in graphs of permutation polyhedra. *Cybernetics and Systems Analysis*. 46(1): pp.7-13 (2010)
36. Kolechkina, L. M., Dvirna, O. A.: Solving extremum problems with linear fractional objective functions on the combinatorial configuration of permutations under multicrteriality. *Cybernetics and Systems Analysis*. 53(4): pp. 590-599 (2017)

37. Kolehckina, L. M., Dvirna, O. A., Nagornaya, A. N.: Modified coordinate method to solve multicriteria optimization problems on combinatorial configurations. *Cybernetics and Systems Analysis*. 50(4): pp. 620-626 (2014)
38. Kolehckina, L., Pichugina, O.: Multiobjective Optimization on Permutations with Applications, In *DEStech Transactions on Computer Science and Engineering*, 2018 IX International Conference on Optimization and Applications (OPTIMA 2018) (Supplementary Volume), pp. 61-75 (2018)